



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

escaped. The fungi peculiar to these galls belong to the genus *Macrophoma*, and are not referable to the species of *Phoma* that inhabit the same hosts.

In a second paper<sup>22</sup> NEGER treats the fungi associated with certain wood-boring beetles, *Xyloterus dispar*, *X. lineatus*, and *Hylecoetus dermestoides*, which form ambrosia upon the walls of their tunnels. The fungus related to *Hylecoetus* is probably a species of *Endomyces*; the other two are closely similar, but not identical, and are not identifiable. The species of *Ceratostomella*, which NEGER formerly mistook for ambrosia fungi in this case, are merely weeds in the fungus garden (as are also yeasts and bacteria), which have no part in producing the edible cells. The larvae of these beetles have thus, in their mouths almost, nutritious food abstracted by the vegetative mycelium from the more distant wood cells, instead of the relatively poor food, the wood itself; moreover, the borings are confined to the sap-wood, where the fungi find appropriate conditions for growth.—C. R. B.

**Ovule and ovulate flower of *Juglans*.**—BENSON and WELSFORD<sup>23</sup> have investigated the ovule and ovulate flower of *Juglans* in reference to the discordant results obtained by VAN TIEGHEM (1869) and NICOLOFF (1905). The "allied genera" examined for comparative study were *Myrica*, *Carpinus*, *Morus*, *Urtica*, and *Rheum*. In brief, it may be said that the account of VAN TIEGHEM was confirmed in all particulars. Interesting phases of "reduction" exhibited by the flowers of *Juglans regia* are as follows: (1) the origin of a dimerous condition from a trimerous, (2) barren placentae with a vascular supply, (3) one mode of the phylogenetic origin of the orthotropous basal ovule from an anatropous parietal type.

More extended conclusions deal with the so-called epigyny of the group considered, and with the ovule in angiosperms. The investigators find in the group "no trace of that form of epigyny which is brought about by the concavity of the axis and sinking and inclusion of the ovary within it," which description hardly applies to epigyny anywhere. It is concluded that the described epigyny of *Amentiferae* need not be regarded as an advanced character, and that the term had better be avoided. The conclusions as to the ovule of angiosperms are: (1) it is appendicular, (2) it is phylogenetically provided with a dual integument, and (3) the vascular supply may be compared with that of the outer integument or "cupule" of *Lagenostoma*.—J. M. C.

**Aridity and evolution.**—Of the external factors which have influenced or caused the evolution of the plant kingdom, MACDOUGAL<sup>24</sup> places much stress

---

<sup>22</sup> NEGER, F. W., *Ambrosiapilze* II. Ber. Deutsch. Bot. Gesells. 27:372-389. pl. 17. figs. 2. 1909.

<sup>23</sup> BENSON, M., AND WELSFORD, E. J., The morphology of the ovule and female flower of *Juglans regia* and of a few allied genera. Annals of Botany 23:623-633. figs. 8. 1909.

<sup>24</sup> MACDOUGAL, D. T., Influence of aridity upon the evolutionary development of plants. Plant World 12:217-231. 1909.